



EUI WORKING PAPERS IN ECONOMICS

EUI Working Paper ECO No. 95/19

Learning, Matching and Aggregation

ED HOPKINS

WP
330
EUR

European University Institute, Florence

EUROPEAN UNIVERSITY INSTITUTE



3 0001 0021 7627 1



EUROPEAN UNIVERSITY INSTITUTE, FLORENCE

ECONOMICS DEPARTMENT

EUI Working Paper ECO No. 95/19

Learning, Matching and Aggregation

ED HOPKINS

WP 330
EUR



BADIA FIESOLANA, SAN DOMENICO (FI)

**All rights reserved.
No part of this paper may be reproduced in any form
without permission of the author.**

**© Ed Hopkins
Printed in Italy in June 1995
European University Institute
Badia Fiesolana
I - 50016 San Domenico (FI)
Italy**

Learning, Matching and Aggregation

Ed Hopkins*

European University Institute

Florence 50016, Italy

hopkins@iuecalc.iue.it

May, 1995

Abstract

Fictitious play and “gradient” learning are examined in the context of a large population where agents are repeatedly randomly matched. We show that the aggregation of this learning behaviour can be qualitatively different from learning at the level of the individual. This aggregate dynamic belongs to the same class of simply defined dynamic as do several formulations of evolutionary dynamics. We obtain sufficient conditions for convergence and divergence which are valid for the whole class of dynamics. These results are therefore robust to most specifications of adaptive behaviour.

Journal of Economic Literature classification numbers: C72, D83.

Keywords: Games, Fictitious Play, Learning, Evolution.

*I would like to thank all the people at GREQAM, Marseille, for their hospitality while writing this paper. Alan Kirman, Tilman Börgers and Dan Friedman have given many helpful comments. I would like to thank Josef Hofbauer for pointing out an error in my first attempt to prove Proposition 5

Learning, Motivation and Assessment

John Hattie
University of Melbourne
Australia

Learning is the process of acquiring knowledge, skills, and attitudes through experience, study, or instruction. Motivation is the internal drive that influences the direction, intensity, and persistence of learning. Assessment is the process of gathering information about a learner's knowledge, skills, and attitudes to evaluate their learning progress and inform instruction.

1 Introduction

There has been an increasing interest in using evolutionary models to explain social phenomena, in particular, the evolution of conventions. A common assumption is that there is a large population where individuals are repeatedly randomly matched to play a game. This paper addresses two resulting issues. First, how do learning rules defined for individuals aggregate across such a population. Second, evolutionary models have not achieved universal acceptance. There has been some skepticism as to the degree to which evolutionary dynamics are relevant to economic situations. In an evolutionary system, nature chooses the individuals who embody superior strategies. In human society, individuals learn: they choose strategies that seem superior. There is no certainty that the dynamics generated by the two different processes are identical.

There are a number of potential responses. One adopted by Binmore and Samuelson (1994) is to devise a learning scheme which approximates the dynamics generated by evolution. Thus the results of evolutionary game theory could be recreated by learning. Another is to generalise the evolutionary dynamics by abandoning particular functional forms and looking at wide classes of dynamics which satisfy “monotonicity” or “order compatibility” (Nachbar, 1990; Friedman, 1991; Kandori et al., 1993). The hope is that even if learning behaviour is not identical to evolution, it is sufficiently similar to fall within these wider categories. However, in this paper, a different approach is taken. Rather than designing learning models to suit our purposes, we examine two existing models of learning behaviour current in the literature. This is done in the context of a large random-mixing population.

The question of aggregation of learning behaviour is of interest in its own right. As can be seen in for example, Crawford (1989) or Canning (1992), learning behaviour aggregated across a large population can be qualitatively different from behaviour at the level of the individual. Indeed, we show that aggregation can solve many of the problems encountered in existing learning models. Secondly, the resultant dynamics

are not in general identical to evolutionary dynamics on a similarly defined population. They may not even satisfy monotonicity. However, they all belong to a class of dynamics which for reasons that will become apparent we will call “positive definite”, and share much of their qualitative behaviour.

Fictitious play, our first learning model, was in fact introduced as a means of calculating Nash equilibrium, or in the terminology of the time in order to “solve” games (Brown, 1951; Robinson, 1951). Play was “fictitious” in that it was assumed to be a purely mental process by which agents would decide on a strategy. The fictitious play algorithm selects a pure strategy that is a best reply to the average past play of opponents. One can interpret this as though each player uses past play as a prediction of opponents’ current actions. This is, of course, in the spirit of the adjustment process first suggested by Cournot in the 19th century. While it might not be clear *a priori* where such a naive form of behaviour might lead, in fact, it has been shown, for example, that the empirical frequencies of strategies played approaches a Nash equilibrium profile in zero-sum games (Robinson, 1951) and in all 2×2 games (Miyasawa, 1961).

More recently, fictitious play has again attracted interest, this time as a means of modelling learning¹. This, however, is an interpretation that is problematic. The positive results noted above are qualified by the realisation that convergence of fictitious play is not necessarily consistent with the idea of players “learning” an equilibrium. Convergence to a pure strategy equilibrium is relatively straightforward: after a certain time, each player will keep to a single pure strategy. However, as Young (1993), Fudenberg and Kreps (1993), Jordan (1993) all note, convergence in empirical frequencies to a mixed Nash equilibrium may only entail that play passes through a deterministic cycle (of increasing length) through the strategies in its support. In one sense, players’ “beliefs” converge,

¹Some of the many to have considered fictitious play or similar processes are Canning (1992), Fudenberg and Kreps (1993), Jordan (1993), Milgrom and Roberts (1991), Monderer and Shapley (1993), Young (1993).

even if their actions do not, in that in the limit they will be indifferent between the different strategies in the support of the Nash equilibrium. However, if players' beliefs are predictions of their opponents' play, while correct on average, they are consistently incorrect for individual rounds of play. Implicit in fictitious play is also a strong degree of myopia. In choosing strategies, players take no account of the fact that opponents are also learning. Similarly, if as noted above, play converges to a cycle, players do not respond to the correlated nature of play. Finally, apart from the case of zero-sum games, there is no easy method of determining whether fictitious play converges.

There are other models of learning in games. We can identify a class of learning rules as being based on gradient-algorithms. The behaviour postulated is perhaps even more naive than under fictitious play², indeed, these models were first developed by psychologists and animal-behaviourists for non-strategic settings. More recently they have been applied to game-theory by Harley (1982), Crawford (1985; 1989), Börgers and Sarin (1993), Roth and Erev (1995). Unlike fictitious play-like processes agents do not play a single pure strategy which is a best-reply, agents play a mixed strategy. If a strategy is successful the probability assigned to it is increased, or in the terminology of psychologists, the "behaviour is reinforced". Thus such models are sometimes called "learning by reinforcement" or "stimulus learning". As these models' other name "gradient" suggests, behaviour is meant to climb toward higher payoffs. Adjustment is therefore slower and smoother than under fictitious play. However, the results obtained are not notably more positive. Crawford (1985) showing for example that all mixed strategy equilibria are unstable.

Aggregation can help with these problems. Fudenberg and Kreps (1993) in fact propose the idea of a random-mixing population of players as a justification for the myopia of fictitious play-like learning processes. If there is sufficient anonymity such that each player cannot identify

²There are other models not considered here such as the more sophisticated Bayesian learning of Kalai and Lehrer (1993).

his opponent and sufficient mixing, each player has a sequence of different opponents, then players may have little incentive to develop more sophisticated strategies. A population of players also offers a different interpretation of mixed-strategy equilibrium. The distribution of strategies in the population as a whole mimics a mixed-strategy profile. This is an equilibrium concept familiar from evolutionary game theory. This type of mixed equilibrium can be stable under either fictitious play or gradient learning.

The main contribution of this paper is to demonstrate that it is possible to obtain precise results on the aggregation of learning behaviour and that furthermore, that the aggregate dynamics thereby obtained are qualitatively very similar to evolutionary dynamics. In fact, we show that the replicator dynamics, in both pure and mixed strategy forms, the aggregate dynamics generated by fictitious play, and also the aggregate dynamics generated by gradient learning, all belong to a simply-defined class of dynamics. We then show that for all of this class that regular Evolutionary Stable Strategies (ESSs) are asymptotically stable in a deterministic context and that a similar result holds for time averages in a stochastic context. Thus we show that refinements to Nash equilibrium based on evolutionary considerations are relevant also for learning models. Secondly, unlike existing models of learning in large populations, such as Canning (1992) and Fudenberg and Levine (1993), explicit results on the stability of particular equilibria are obtained. Perhaps most importantly we obtain results which are robust to different specifications of learning rules or evolutionary dynamics. Hence we can hope that these results have some predictive power.

2 Learning and Evolutionary Dynamics

We will examine learning in the context of two-player normal-form games, $G = (\{1, 2\}, I, J, A, B)$. I is a set of n strategies available to player 1, J a set of m strategies for player 2. Payoffs are determined by A , a

$n \times m$ matrix of payoffs, and B , which is $m \times n$. A has typical element a_{ij} , which is the payoff an agent receives when playing strategy i against an opponent playing strategy j . However, we will largely be dealing with games that are “symmetric” in the evolutionary sense, that is, games for which $A = B^3$. Generalisations to the asymmetric case are straightforward. We will often be dealing with a population of players, each playing a single pure strategy. In this case, the distribution of strategies within the population will be described by a vector $\mathbf{x} \in S_n = \{\mathbf{x} = (x_1, \dots, x_n) \in \mathbf{R}^n : \sum x_i = 1, x_i \geq 0 \text{ for } i = 1, \dots, n\}$. As, in this paper, vectors will be treated ambiguously as either rows or columns, to avoid any further confusion, the inner product will be carefully distinguished by the symbol “ \cdot ”, that is, the result of $\mathbf{x} \cdot \mathbf{x}$ is a scalar.

We follow Shapley (1964) and implement the fictitious play algorithm in the following way. A player places a weight on each of her strategies (we can think of these as beliefs as to the relative effectiveness of the different strategies) which we can represent as a vector $\mathbf{w} = (w_1, w_2, \dots, w_n)$ and at any given time plays the strategy which is given the highest weight. Each player updates these weights after each round of play so that if her opponent played strategy j ,

$$w_i(t+1) = w_i(t) + a_{ij} \text{ for } i = 1, \dots, n. \quad (1)$$

Players can also be modelled as maintaining a vector of relative frequencies of opponents’ past play (as in Fudenberg and Kreps, 1993; Young, 1993). They then choose strategies that maximise expected payoffs as though this vector represented the current (mixed) strategy of their opponents. The two methods are entirely equivalent. Note that the weights here are (less initial values) simply the relative frequencies multiplied by payoffs.

Up to now we have contrasted learning and evolution purely on the basis of their origins, one being a social, the other a natural process.

³And all players are drawn from the same population. For a fuller discussion of the difference between symmetric and asymmetric contests see van Damme (1991) or Hofbauer and Sigmund (1988).




Figure 1: Dynamics: (a) best response (b) replicator dynamics

However, they are also often modelled in contrasting fashion. Fictitious play and Cournotian dynamics both assume that agents play some kind of best response. This can involve discontinuous jumps in play. Taking as an example the following game which is variously known as “chicken”, “hawk-dove” or “battle of the sexes”,

$$A = B = \begin{array}{|c|c|} \hline 0 & a \\ \hline 1-a & 0 \\ \hline \end{array} \quad 1 > a > 0, \quad (2)$$

Figure 1a gives the simple best-reply function for (2), where each agent in a large population plays a best-reply to the current distribution of strategies⁴. Here x represents the proportion of the population playing the first strategy. If x is greater than (respectively less than) a , then the whole population switches to strategy 2 (strategy 1). Hence, there is a discontinuity at the point ($x = a$) where the players are indifferent between their two strategies (there is no particular consensus in the literature about how players should behave when indifferent between two or more strategies).

In contrast, the evolutionary *replicator dynamics*, whether in continuous or discrete time, are derived on the basis that the proportional

⁴This is a dynamic as used by, for example Kandori, Mailath and Rob (1994). This is fictitious play with a one-period memory.

rate of growth of each strategy is equal to the difference between its payoff $(A\mathbf{x})_i$ (the i th element of the vector in parentheses) and the average payoff in the population⁵ $\mathbf{x} \cdot A\mathbf{x}$. D is a positive constant.

$$\dot{x}_i = x_i[(A\mathbf{x})_i - \mathbf{x} \cdot A\mathbf{x}] \text{ or } x_i(t+1) = x_i(t) \frac{(A\mathbf{x})_i + D}{\mathbf{x} \cdot A\mathbf{x} + D} \quad (3)$$

Clearly, both dynamics are continuous, the system moving smoothly toward the strategies earning the highest payoff. The replicator dynamic (in discrete time) for the game (2) is drawn in Figure 1b. The interior mixed equilibrium is a global attractor, the pure equilibria at $x = 0, 1$ being unstable.

Important in evolutionary theory is the idea of an Evolutionary Stable Strategy, that is, “a strategy such that, if all members of a population adopt it, then no mutant strategy could invade the population under the influence of natural selection.” (Maynard Smith, 1982, p10). For a large random matching population the conditions are

Definition: An *Evolutionary Stable Strategy* (ESS) is a strategy profile \mathbf{q} that satisfies the Nash equilibrium condition

$$\mathbf{q} \cdot A\mathbf{q} \geq \mathbf{x} \cdot A\mathbf{q} \quad (4)$$

for all $\mathbf{x} \in S_n$ and for all \mathbf{x} such that equality holds in (4), \mathbf{q} must also satisfy the stability condition

$$\mathbf{q} \cdot A\mathbf{x} > \mathbf{x} \cdot A\mathbf{x} \quad (5)$$

The first condition states that to be an ESS, a strategy must be a best-reply to itself. Were it not so, a population playing that strategy could easily be invaded by agents playing the best reply. The second condition demands that if there are a number of alternative best replies,

⁵In a biological context, this arises from relative reproductive success (see Hofbauer and Sigmund, 1988) but may also be an appropriate assumption in modelling learning in a human population (for example, Binmore and Samuelson, 1994).

than the ESS must be better against them than they are against themselves. Thus if a mutant strategy which was an alternative best reply were to enter the population, those agents playing it would on average have a lower payoff than those playing the ESS and therefore would not grow in number.

There is a strong connection between stability under evolutionary dynamics and the static concept of ESS.

Proposition 1 *Every ESS is an asymptotically stable equilibrium for the continuous time replicator dynamics but the converse is not true. That is, there are asymptotically stable states for the replicator dynamics which are not ESSs.*

Proof: See, for example, van Damme (1991, Theorem 9.4.8). \square

Fictitious play can also converge on the mixed equilibrium of (2), but in a rather different manner. Setting $a = 0.5$, imagine two players both with initial weights of $(0.25, 0)$. That is, they both prefer their first strategy for the first round of play. Both consequently receive a payoff of 0. Each player observes which strategy the opponent chose. They then update the weights/beliefs according to the payoffs that they would receive against that strategy. Thus according to (1), weights now stand at $(0.25, 0.5)$. They now both prefer the second strategy. One can infer that player 1 believes that her opponent will continue to play her first strategy, and likewise for player 2. After the second round of play, in which again both players receive 0, the vectors stand at $(0.75, 0.5)$. It can be shown that, firstly, that the players continually miscoordinate, always receiving a payoff of 0, and that, secondly, in the limit, both play their first strategy with relative frequency 0.5, and their second with frequency 0.5. This corresponds to the mixed strategy equilibrium of (2). However, the players' behaviour seems to correspond only tangentially with the idea of a mixed-strategy equilibrium.

The concept of a mixed strategy equilibrium in use in evolutionary game theory seems more intuitive. It is also an average but not across

time but across the differing behaviour of a large population: the aggregate strategy distribution is a mixed strategy equilibrium. One might hope that if each individual used a learning rule that like the replicator dynamics was a continuous function of payoffs, similarly well-behaved results could be obtained. However, Crawford (1985; 1989) demonstrates that in fact mixed strategy equilibria, and hence many ESSs, are not stable for a model of this kind. However, while these results are correct, they do not tell the whole story in the context of a random-mixing population. The mixed strategy of individuals will not approach the equilibrium of the two player game, nonetheless, we are able to prove convergence for the mean strategy in the population for all regular ESSs.

What we are going to show is that with a large population of players who are continually randomly matched, this type of outcome is possible even under fictitious play. This does not follow automatically from aggregation. In particular, if all players in the population have the same initial beliefs, the time path for the evolution of strategies will be the same as for fictitious play with two players⁶. Imagine in the above example, there were an entire population of players with initial weights of $(0.25, 0)$. No matter with whom they are matched they will meet an opponent playing, strategy 1. Hence, all players will update their beliefs at the same rate, and the same cycle is reproduced. However, this is only possible given the concentration of the population on a single point. If instead there is a non-degenerate distribution of weights across the population, it may be that not all the population will change strategy at once.

Imagine now that the players have initial weights or beliefs $(b, 0)$ where b is uniformly distributed on $[0, 1]$. Only those in the population with $b \leq 0.5$, that is half the population, will change strategy after the first round of play. In fact, we have arrived immediately at the population state equivalent to the mixed strategy equilibrium with half the population playing each strategy. It is easy to check that under random matching, in such a state, there is no expected change in each

⁶A fact which Fudenberg and Kreps (1993) exploit. They do not consider the case where, within a population of players, individuals possess differing beliefs.

player's strategy. In this case, aggregation has had a smoothing effect because there was sufficient heterogeneity in the population. We will go on to make a somewhat more precise statement about convergence of fictitious play in a random matching environment. A necessary first step is to consider the modelling of random matching itself in more detail.

3 Matching Schemes

Any study of the recent literature on learning and evolution will reveal, firstly, that random matching within a large population of players is a common assumption, and secondly, that there are several ways of modelling such interaction. This diversity is in fact important both in terms of what it implies for theoretical results and in what cases are such results applicable. For example, there are some economic or social situations where random matching might seem a reasonable approximation of actual interaction, others where it will not. Only in some cases will agents be able to obtain information about the result of matches in which they were not involved, and so on.

Fudenberg and Kreps (1993) suggest three alternative schemes. Assuming a large population of potential players (they suggest 5000 as a reasonable number), they propose the following:

“Story 1. At each date t , one group of players is selected to play the game...They do so and their actions are revealed to all the potential players. Those who play at date t are then returned to the pool of potential players.

Story 2. At each date t there is a random matching of all the players, so that each player is assigned to a group with whom the game is played. At the end of the period, it is reported to all how the entire population played....The play of any particular player is never revealed.

Story 3. At each date t there is a random matching of the players, and each group plays the game. Each player recalls

at date t what happened in the previous encounters in which he was involved, without knowing anything about the identity or experiences of his current rivals.”

Fudenberg and Kreps (1993, p333)

It is worth drawing out the implications of these different matching schemes. We will argue that Story 3 is the most appropriate basis for a model of fictitious play in a population of players. It is the “classic” scheme assumed as a basis for the replicator dynamics. The population is assumed to be infinite and hence, despite random matching, the dynamics are deterministic (this has been rigorously analysed by Boylan, 1992). It is also decentralised and does not require, as do Stories 1 and 2, any public announcements of results by some auctioneer-like figure. However, there are other procedures similar to Story 2 which do not require such a mechanism. These include,

Story 2a. In *each* round⁷, the players are matched according to Story 1 or Story 3 an infinite number of times.

Story 2b. In *each* round there is a “round-robin” tournament, where each player meets each of his potential opponents exactly once.

Stories 2a and 2b have been used in the learning literature principally for reasons of tractability⁸. They ensure a deterministic result to the matching procedure even when population size is finite. The infinite number of matchings in Story 2a, by the law of large numbers, ensures that a proportion equal to the actual frequency over the whole population of opponents playing each strategy will be drawn to play. What Stories 2, 2a and 2b have in common is that all players know the exact distribution of strategies in the population when choosing their next strategy.

⁷The “round” is the time-unit of, in evolutionary models, reproduction, in learning models, decision. That is, strategy frequencies are constant within a round, even if the round contains many matches.

⁸See for example, Kandori et al. (1993), Binmore and Samuelson (1994).

There is little room for the diversity of beliefs one might expect in a large population.

In contrast, under Story 3, as the overall distribution of strategies is not known, it makes more sense to use past matches to estimate the current distribution. Furthermore, depending upon with which opponent they are matched, different players will receive different impressions about the frequency of strategies in the population of opponents. Under Story 3, if the population is finite, even if players use a deterministic rule to choose their strategy, such as the fictitious play algorithm, the evolution of the aggregate strategy distribution is stochastic. We therefore develop two models, one deterministic and one stochastic, which describe the aggregate behaviour of the population under the different schemes.

4 Population Fictitious Play

The next stage is to examine population fictitious play (PFP) where learning takes place in a large random-mixing population. We will deal both with the case where the population is large but finite, and with the case where the population is taken to be a continuum of non-atomic agents (an assumption familiar from evolutionary game theory). While the beliefs of a given individual can be represented by point, the beliefs of the population will be represented by a distribution over the same space. We investigate how the distribution of beliefs, and therefore how the distribution of strategies, changes over time. It will help to create some new variables. Let $p_{ij} = w_j - w_i$, $j \neq i$. Thus, \mathbf{p}_i is a vector of length $n - 1$. We will use this to work in \mathbf{R}^{n-1} instead of \mathbf{R}^n . For example, if a player has to choose between two strategies, we can summarise her beliefs by the variable p_{12} . If $p_{12} < 0$ she prefers her first strategy, if $p_{12} > 0$ her second, and if $p_{12} = 0$ she is indifferent. A player's decision rule or reaction function can then be considered as a mapping from the space of beliefs to strategies, i.e. $\mathbf{R}^{n-1} \rightarrow S_n$, that is, the n -simplex. This mapping will not, in general be continuous for individual players:

the fictitious play assumption limits players to pure strategies. See also Figure 1a.

Let F_i be the population distribution function of \mathbf{p}_i over \mathbf{R}^{n-1} . Agents will play a strategy if it is the strategy given the highest weight in their beliefs. In other words, the beliefs of those playing strategy i must be in $E_i = \{\mathbf{p}_i \in \mathbf{R}^{n-1} : p_{ij} \leq 0, \forall j \neq i\}$. What if agents are indifferent between two or more strategies, that is, if their beliefs, for some j are such that $p_{ij} = 0$? One way to finesse this problem is to assume that payoffs and/or initial beliefs are given by irrational numbers. Another method is to assume that beliefs are given by a continuous distribution on \mathbf{R}^{n-1} , (F is absolutely continuous). If this is case then, if the proportions of the population playing each of the n strategies is given by the vector $\mathbf{x} \in S_n$, $x_i = F_i(\mathbf{0})$, where $\mathbf{0}$ is a vector of zeros of length $n - 1$. For example, if all agents have the beliefs $p_{ij} < 0 \forall j$ then $x_i = F_i(\mathbf{0}) = 1$.

At the basis of the deterministic model of PFP is the assumption that agents update their beliefs as if they knew $\mathbf{x} \in S_n$, the true current distribution of strategies in the population. This could be supported by Story 3 in an infinite population or by Story 2 in a finite or infinite population. We are, however, going to treat each x_i as a continuous variable and assume that the probability of meeting an opponent playing strategy i is x_i .⁹ For example, over a period of length Δt , each agent is matched within a single large population. If this matching is repeated an arbitrarily large number of times in each period (Story 2a), each agent will meet a proportion x_i of opponents playing strategy i . According to (1), which describes the fictitious play algorithm, we have for each agent

$$\mathbf{w}(t + \Delta t) = \mathbf{w}(t) + A\mathbf{x}. \quad (6)$$

Similarly we can derive a system of difference equations for \mathbf{p} , the vector of the agent's beliefs,

$$\mathbf{p}_i(t + \Delta t) = \Gamma(\mathbf{p}_i, \mathbf{x}) = \mathbf{p}_i(t) + [(A\mathbf{x})_{j \neq i} - (A\mathbf{x})_i], \quad (7)$$

⁹These are both approximations if the population is finite. We treat finite populations with greater accuracy in Section 7.




Figure 2: Change in the distribution of beliefs

where $(A\mathbf{x})_{j \neq i}$ is a vector of length $n-1$, constructed of all the elements of $A\mathbf{x}$ except $(A\mathbf{x})_i$. We will be interested in the properties of the inverse of the function Γ with respect to \mathbf{p}_i to be written $\Gamma^{-1}(\mathbf{p}_i)$. Given that $\Gamma(\cdot)$ is a simple linear function the existence of Γ^{-1} is therefore guaranteed. In fact, we have

$$\Gamma^{-1}(\mathbf{p}_i) = \mathbf{p}_i(t) + [(A\mathbf{x})_i - (A\mathbf{x})_{j \neq i}] \quad (8)$$

To illustrate the properties of the deterministic model with a simple example, we consider 2×2 symmetric games, that is, games where every player must choose between the same two strategies. Let $F_t(p)$ be the cumulative distribution of $p = p_{12} = -p_{21}$ on \mathbf{R} . This distribution of beliefs determines the distribution of strategies. As the t subscript indicates, this distribution will change endogenously over time, as the beliefs of each agent are updated according to (7). This is shown in Figure 2, (in the figure, a density function $f = dF/dp$ is assumed; its existence is

not necessary to the analysis of this section). In particular,

$$\begin{aligned} \Gamma^{-1}(p) > p : F_{t+\Delta t}(p) &= F_t(p) + \int_p^{\Gamma^{-1}(p)} dF = F_t(\Gamma^{-1}(p)) \\ \Gamma^{-1}(p) < p : F_{t+\Delta t}(p) &= F_t(p) - \int_{\Gamma^{-1}(p)}^p dF = F_t(\Gamma^{-1}(p)) \end{aligned} \quad (9)$$

Any agents possessing beliefs equal to $\Gamma^{-1}(0)$ will update their beliefs to p_0 . If $\Gamma^{-1}(0) > 0$, as is the case in Figure 2, $F(0)$ will increase by the proportion of agents who possessed beliefs on the interval $[0, \Gamma^{-1}(0)]$. The linear nature of (7) implies that the whole distribution simply shifts to the left or to the right. This in turn will have an effect on the distribution of strategies. For example, an agent whose beliefs change from $p = 1$ to $p = -1$ will change from her second to her first strategy. By definition, if F is continuous at $p = 0$, that is, there is no mass of agents indifferent between strategies, $x_1 = F(0)$ and hence

$$x_1(t + \Delta t) = F_t(\Gamma^{-1}(0)) = F_t([(A\mathbf{x})_1 - (A\mathbf{x})_2]). \quad (10)$$

That is, in Figure 2, x_1 increases by an amount equal to the shaded area. It is not difficult to extend this analysis to games of n strategies. In a time interval of length Δt , the change in x_i is given by

$$x_i(t + \Delta t) = F_t(\Gamma^{-1}(\mathbf{0})) = F_{it}([(A\mathbf{x})_i - (A\mathbf{x})_{j \neq i}]), \quad (11)$$

where F_i is the joint cumulative distribution function of \mathbf{p}_i on \mathbf{R}^{n-1} . Clearly, if a strategy i currently has a higher expected payoff than any other strategy, then the proportion of the population playing that strategy x_i is increasing. We can state that more formally as:

Lemma 1 *If $(A\mathbf{x})_i > (A\mathbf{x})_j \forall j \neq i$ then p_{ij} is strictly decreasing at a rate bounded away from zero $\forall j \neq i$ and x_i is increasing. If $(A\mathbf{x})_i = (A\mathbf{x})_j \forall j$ then $p_{ij} \forall j \neq i$, and $x_i \forall i$, are constant.*

While the state variable of the PFP process is the distribution of agents' beliefs, our main focus of interest is the distribution of strategies. We therefore define a fixed point for the PFP process as a population strategy profile which is unchanging under the dynamic specified by (7),

even though beliefs may continue to change. We find a one-to-one correspondence between fixed points and strategy distributions that are Nash equilibria of the game. Mixed strategies are supported by the appropriate distribution of pure strategies across the population.

Proposition 2 *A strategy profile \mathbf{q} in the simplex S_n is a fixed point for the deterministic PFP dynamic if and only if it is a Nash equilibrium.*

Proof: We can start by observing that if \mathbf{q} is a Nash equilibrium then from (4) above, if $I_0 \subseteq I$ is the set of strategies in the support of \mathbf{q} , then

$$\forall i, j \in I_0 \quad (A\mathbf{q})_i = (A\mathbf{q})_j \geq (A\mathbf{q})_k \quad \forall k \notin I_0 \quad (12)$$

(a) **If.** If an agent plays i , she must prefer it. That is, $w_i \geq w_j \quad \forall j$. From Lemma 1 and (12), no agent will change preference either between the strategies in the support of \mathbf{q} or toward any other strategy.

(b) **Only if.** Let \mathbf{q} now denote a rest point which is not a Nash equilibrium. Let $I_0 \subseteq I$ be the set of strategies in its support. If \mathbf{q} is not a Nash equilibrium then there must be a set of strategies I_k such that $\exists i \in I_0 \quad (A\mathbf{q})_i < (A\mathbf{q})_k \quad \forall k \in I_k$. From (7), for each agent playing strategy i , $w_i - w_k$ must be decreasing at a constant rate as long as the system is at \mathbf{q} . Within finite time, a positive measure of agents playing i must switch to a strategy in I_k . Hence the system is no longer at \mathbf{q} . \square

The following propositions are also immediate consequents.

Proposition 3 *All pure strict Nash equilibria are asymptotically stable.*

Proof: A pure strict Nash equilibrium is a state $\mathbf{q} \in S_n$ with one strategy i in its support such that there exists an open ball B with centre \mathbf{q} such that in $B \cap S_n$, $(A\mathbf{x})_i > (A\mathbf{x})_j \quad \forall j \neq i$. Clearly, if the system enters B by the previous Lemma it cannot leave. While in B , for all agents, each $p_{ij} \quad \forall j \neq i$ is decreasing at a non-vanishing rate. In finite time, all agents play i . \square

Proposition 4 *All strictly dominated strategies have zero population share in finite time.*

Proof: This follows immediately from Lemma 1. □

These results are hardly surprising given that we have a population of agents that play only best replies, but they are sufficient to show convergence for many games. However, because mixed strategy equilibria are never strict, to deal with them we will need to change our approach.

5 Positive Definite Dynamics

We will now modify our existing model in two important ways. First, we will move from discrete to continuous time. This is not a neutral step. Our defence is that a discrete time model implies that all players are matched, and hence update their behaviour, simultaneously, a degree of coordination unlikely in a large population. Second, it is necessary to impose additional assumptions to ensure that the distribution of beliefs is continuous. For example, if there were mass points, discontinuous jumps in the value of \mathbf{x} would be possible as positive measures of players switched beliefs. As we have seen the deterministic cycles of normal fictitious play are possible even in the large population model, but only with extreme restrictions on initial beliefs. Indeed, any perturbation to the distribution of beliefs will change the dynamic behaviour substantially.

Zeeman (1981) faced a similar problem in modelling mixed-strategy evolutionary dynamics. We follow the same strategy of assuming that the distributions we consider are subject to noise. For Zeeman, who was considering a biological model this was caused by mutations. Here, we can either assume that players make idiosyncratic, independently distributed mistakes in updating their beliefs, or, in the spirit of purification (see also Fudenberg and Kreps, 1993), we can imagine that each individual payoffs are subject to idiosyncratic shocks. More formally, we imagine a

once-off shock of the form:

$$\mathbf{w}(t + \Delta t) = \mathbf{w}(t) + \eta, \quad (13)$$

where η is a vector of normally-distributed independent random variables each with zero mean and finite variance. This would rule out the possibility of mass points of agents holding exactly the same beliefs. For example, in the two strategy case, if $p = -1$ for all agents, that is, they all prefer their first strategy, with the addition of the noise, beliefs would instead be normally distributed with mean -1. We can choose the variance of η sufficiently small such that the new distribution approximates the old arbitrarily closely. Indeed, as Zeeman notes, distributions which satisfy our conditions are open dense in the set of all distributions. We state these conditions in more detail:

Assumption of Continuity: the distribution of beliefs is such that F_i is absolutely continuous with respect to \mathbf{p}_i . There exist continuously differentiable density functions $f_{ij} = f_{ji} = dF_i/dp_{ij}$ on \mathbf{R}^{n-1} such that $f_{ij} > 0$ everywhere on \mathbf{R}^{n-1} .

The last inequality in turn implies that $x_i(t) > 0 \forall i, t$. However, it is possible for the system to approach the boundary of the simplex asymptotically. Consider the case where there is a single strictly dominant strategy i . In the previous section, we saw that, without noise, within a finite time only that strategy would be played. Here, the noise means that some agents will always prefer other strategies, but over time the numbers doing so will drop away to zero. The reason is that, from (13), we have $E[p_{ij}(t + \Delta t) - p_{ij}(t)] < 0 \forall j \neq i$, the strength of preference for the dominated strategies is always decreasing. The result is that, $\lim_{t \rightarrow \infty} \Pr[w_j + \eta_j > w_i + \eta_i] = 0$. Hence, $\lim_{t \rightarrow \infty} x_j = 0$ and $\lim_{t \rightarrow \infty} F(\mathbf{0}) = 1$.

We are now going to take the continuous time limit. Returning to Figure 2, in discrete time, all agents with beliefs in the interval $[0, \Gamma^{-1}(0)]$ changed strategy. As we will see, moving to continuous time is equivalent

of taking the limit $\Gamma^{-1}(0) \rightarrow 0$. That is, the rate of change at any given point in time is going to depend on the number of agents who are, at that instant, passing from preference of one strategy to preference of another. In other words, the rate of change will be proportional to the density of agents at the point of indifference, in Figure 2, $f(0)$. Subtracting x_i from both sides of (11),

$$x_i(t + \Delta t) - x_i(t) = F_{it}([(\mathbf{Ax})_i - (\mathbf{Ax})_{j \neq i}]) - F_{it}(\mathbf{0}). \quad (14)$$

Given the presence of a random disturbance in (13), the reader may be surprised to see none in the above formula. The errors, however, have been subsumed in the distribution function F_i . Next, we divide through by Δt and take the limit $\Delta t \rightarrow 0$ to obtain

$$\dot{x}_i = \sum_{j \neq i} [(\mathbf{Ax})_i - (\mathbf{Ax})_j] f_{ij}(\mathbf{0}). \quad (15)$$

If there is any doubt about this derivation, note that, it satisfies the chain rule $dF_i/dt = dF_i/d\mathbf{p}_i \cdot d\mathbf{p}_i/dt$, where the last term is the continuous time limit of (8). It is also consistent with the theory of surface integrals which scientists and engineers use to calculate the flow of fluid (in this case, beliefs) across a surface. It will be useful to write (15) in matrix form,

$$\dot{\mathbf{x}} = Q(F(t))\mathbf{Ax}. \quad (16)$$

(For the sake of simplicity, we will often suppress the extra arguments that follow Q). Clearly, (15) is very close to the continuous-time replicator dynamics (3) and the linear dynamics proposed by Friedman (1991),

$$\dot{x}_i = \frac{1}{n} \sum_{j \neq i} [(\mathbf{Ax})_i - (\mathbf{Ax})_j] \quad (17)$$

In particular, if the distribution of beliefs is symmetric, such that $f_{ij} = f_{ik}, \forall j, k$, then the continuous time PFP is identical to the linear dynamics. However, if the distribution is such that $f_{ij} = x_i x_j$, then the replicator dynamics are reproduced. In any case, without placing any restrictions on the shape of the distribution, we have the following results

Lemma 2

1. Every element of Q is continuously differentiable in \mathbf{x} ,
2. $\lim_{x_i \rightarrow 0} Q_{ij} = 0 \forall j$,
3. $Q\mathbf{u} = \mathbf{0}$, where \mathbf{u} denotes the vector $(1, 1, \dots, 1)$,
4. Q is positive semi-definite. That is, if $A\mathbf{x}$ is not a multiple of \mathbf{u} then $A\mathbf{x} \cdot QA\mathbf{x} > 0$,
5. Q is symmetric.

Proof: Q has a diagonal $Q_{ii} = \sum_{j \neq i} f_{ij}$ and off-diagonal $Q_{ij} = Q_{ji} = -f_{ij}$. Satisfaction of Conditions 1 and 2 is guaranteed by the Continuity Assumption. Hence at a vertex of S_n , Q consists of zeros. Clearly $Q\mathbf{u} = \mathbf{u} \cdot Q = \mathbf{0}$. However, $\mathbf{x} \cdot Q\mathbf{x} = \sum_{j \neq i} f_{ij}(x_i - x_j)^2 \geq 0$. \square

Geometrically, the operator Q maps the vector of payoffs $A\mathbf{x}$ from \mathbf{R}^n to the subspace $\mathbf{R}_0^n = \{\mathbf{z} \in \mathbf{R}^n : \mathbf{u} \cdot \mathbf{z} = 0\}$ (if the vector $QA\mathbf{x}$ did not add to zero then \mathbf{x} would cease to add to one). It has nullspace \mathbf{u} . That is, at a mixed Nash equilibrium where payoffs are equal ($A\mathbf{x}$ is a multiple of \mathbf{u}), $\dot{\mathbf{x}} = \mathbf{0}$. For other Nash equilibria, if a strategy j is not in the support of \mathbf{q} , then at \mathbf{q} , $f_{ij} = 0$. Because Q is positive definite the angle between $A\mathbf{x}$ and $QA\mathbf{x}$ is less than 90° . This last property is what Friedman (1991) calls “weak compatibility”.

Definition: Any dynamic of the form $\dot{\mathbf{x}} = QA\mathbf{x}$, where the matrix Q , satisfies the above 5 conditions, we call a *positive definite dynamic*.

We can demonstrate that evolutionary concepts are important in the context of population fictitious play. In particular, we can show that all ESSs are asymptotically stable. First we need a preliminary result,

Lemma 3 Any ESS \mathbf{q} is negative definite with respect to the strategies in its support. That is, $(\mathbf{x} - \mathbf{q}) \cdot A(\mathbf{x} - \mathbf{q}) < 0$ for all \mathbf{x} with the same support as \mathbf{q} (see van Damme, 1991; Theorem 9.2.7).

The following lemma and proposition are based upon work of Hines (1980), Hofbauer and Sigmund (1988) and Zeeman (1981). However, the result obtained here generalises the above results and indeed extends beyond the continuous time PFP process to any dynamics which are symmetric positive definite transformations of the vector of payoffs $A\mathbf{x}$.

Lemma 4 *If A is negative definite when constrained to \mathbf{R}_0^n (that is, $\mathbf{z} \cdot A\mathbf{z} < 0 \forall \mathbf{z} \in \mathbf{R}_0^n$), then QA is a stable matrix (i.e. all its eigenvalues have negative real parts when QA is constrained to \mathbf{R}_0^n).*

Proof: The eigenvalue equation is $QA\mathbf{z} = \lambda\mathbf{z}$ for some $\mathbf{z} \in \mathbf{C}_0^n = \{\mathbf{z} = \mathbf{z}_1 + \mathbf{z}_2i \in \mathbf{C}^n : \mathbf{z}_1, \mathbf{z}_2 \in \mathbf{R}_0^n\}$. We can construct a vector \mathbf{y} such that $\mathbf{z} = Q\mathbf{y}$, where $\mathbf{z} \in \mathbf{C}_0^n$. By the symmetry of Q , we have $\mathbf{y}^c \cdot Q = \mathbf{z}^c$ where \mathbf{z}^c is the conjugate of the complex vector \mathbf{z} . This gives us

$$\mathbf{y}^c QA\mathbf{z} = \mathbf{z}^c \cdot A\mathbf{z} = \lambda\mathbf{y}^c \cdot \mathbf{z} = \lambda\mathbf{y}^c \cdot Q\mathbf{y} \quad (18)$$

As Q is symmetric positive definite, $\mathbf{y}^c \cdot Q\mathbf{y}$ is real and positive. The real part of $\mathbf{z}^c \cdot A\mathbf{z}$ is negative, hence the real part of λ is negative. Since all its eigenvalues are negative or have negative real part for eigenvectors in \mathbf{R}_0^n , QA is a stable matrix on that space. \square

A strategy profile \mathbf{q} is a **regular ESS** if it is an ESS that satisfies the additional requirement that all strategies that are a best reply to \mathbf{q} are in its support. We are now able to prove

Proposition 5 *All regular ESSs are asymptotically stable for any positive definite dynamic.*

Proof: Let \mathbf{q} be a fully mixed ESS. Differentiating $Q(\mathbf{x})A\mathbf{x}$ with respect to \mathbf{x} and evaluating at \mathbf{q} , we obtain $Q(\mathbf{q})A + dQ/d\mathbf{x} A\mathbf{q}$. At a Nash equilibrium $QA\mathbf{x} = \mathbf{0}$. It follows that for each x_i , $dQ/dx_i A\mathbf{q} = \mathbf{0}$. Thus the Jacobian of the system at \mathbf{q} is given by $Q(\mathbf{q})A$. By Lemma 4 all its eigenvalues have real part negative.

If a regular ESS \mathbf{q} is on a face $S_q \subset S_n$, that is, $q_i > 0$ if and only if $i \in I_q \subset I$, then it is also asymptotically stable under the continuous time positive definite dynamic. Because it is an ESS, A is a negative definite form on S_q , and so is QA is stable on S_q . It remains to show that the dynamic will approach S_q from the interior of S_n .

We adapt the proof of Zeeman (1981). Define $\lambda = \mathbf{u} \cdot \mathbf{q} \cdot A\mathbf{q} - A\mathbf{q}$. This is a vector whose i th element is zero for $i \in I_q$ and positive for $i \notin I_q$. Hence, we can define the function $\Lambda = \lambda \cdot \mathbf{x} \geq 0$, with equality on S_q , and $\dot{\Lambda} = \lambda \cdot Q\mathbf{A}\mathbf{x}$. We choose an ϵ such that for all \mathbf{x} in some neighbourhood of \mathbf{q} , $\mathbf{x} = \mathbf{q} + \xi$ with $|\xi_i| < \epsilon$, and $|Q_{ij}| < \epsilon$ for $i \notin I_q$ by Conditions 1 and 2 of the definition of a positive definite dynamic. Then

$$\dot{x}_i = \sum_j Q_{ij}(A\mathbf{q})_j + \sum_{j,k} Q_{ij} \dot{A}_{jk} \xi_k$$

Now, if $i \notin I_q$ then the first term of the above is of order ϵ , the second is of the order ϵ^2 . Thus, in the neighbourhood of \mathbf{q} we can approximate $\dot{\Lambda}$ by $\lambda \cdot Q(\mathbf{u} \cdot \mathbf{q} \cdot A\mathbf{q} - \lambda) = -\lambda \cdot Q\lambda < 0$. \square

What is particularly attractive about this result is that to determine stability one no longer has to examine the potentially complicated function $Q(\mathbf{x})$. Instead, one can confine attention to the properties of A alone. For example, for the PFP dynamics it is not necessary to know the shape of the distribution of beliefs. The last two conditions on Q are the substantive ones. Positive definiteness is a necessary condition for stability. (If there is a \mathbf{y} such that $\mathbf{y} \cdot Q\mathbf{y} < 0$, set $\mathbf{z} = Q\mathbf{y}$ and consider (18). If A is negative definite QA must have at least one positive eigenvalue.) It becomes sufficient when combined with symmetry. Why this should lead to asymptotic stability for ESSs can be seen in the traditional economic terms of convexity and concavity. A “positive definite” dynamic is a gradient-climber. The negative definiteness of ESSs of course implies concavity. Any positive definite dynamic will move “uphill” toward the ESS. Condition 1 is the necessary condition for an unique solution to the differential equation (16). Condition 2 ensures that the dynamic remains upon the simplex. Of course, both the replicator dynamics and

Friedman's linear dynamics satisfy these conditions (the latter only on the interior of the simplex).

The importance of symmetry can be illustrated by comparing positive definiteness with the Friedman's (1991) concept of order compatibility or the monotonicity of Nachbar (1990) and Samuelson and Zhang (1992). Monotonicity requires that $\dot{x}_i/x_i > \dot{x}_j/x_j$ iff $(A\mathbf{x})_i > (A\mathbf{x})_j$, and order compatibility, $\dot{x}_i > \dot{x}_j$ iff $(A\mathbf{x})_i > (A\mathbf{x})_j$. It is easy to check that if a dynamic can be written $\dot{\mathbf{x}} = Q(\mathbf{x})A\mathbf{x}$ both monotonicity and order compatibility imply the positive definiteness of Q (as Friedman points out order compatibility implies weak compatibility which is equivalent to positive definiteness). However, monotonicity and order compatibility do not imply symmetry. The existence of asymmetric order-compatible dynamics is what enables Friedman (1991) to demonstrate that ESSs may be unstable under order compatible dynamics. Similarly, there are dynamics which are monotonic but which diverge from ESSs. Conversely, there are positive definite dynamics which are not monotone or order compatible.

In the case of only two strategies, for any such positive definite dynamic, we have

$$\dot{x}_1 = Q_{11}[(A\mathbf{x})_1 - (A\mathbf{x})_2] \tag{19}$$

For 2×2 games, the orbits produced by the positive definite dynamics will, after a suitable rescaling, be identical.

Proposition 6 *For 2×2 games, all positive definite dynamics generate orbits which are identical up to a change in velocity.*

Proof: Continuous dynamical systems are invariant under positive transformations, which represent a change in velocity (see for example, Hofbauer and Sigmund, 1988, p92). By positive definiteness Q_{11} is positive on the relevant state space. \square

6 Mixed Strategy Dynamics

The replicator dynamics do not allow individuals the use of mixed strategies. As van Damme (1991) notes it would be preferable to examine mixed strategy dynamics which permit this possibility. The problem is that they are less tractable than the replicator dynamics which they generalise. In this section, we are able to show that they also fall within the class of positive definite dynamics. Furthermore, we show that the aggregation of gradient learning can be treated in a similar manner.

Zeeman (1981, Section 5) examines the properties of the mixed-strategy replicator dynamics (see also Hines, 1980). The main assumption is that there is an infinite random-mixing (Story 3) population whose individuals play mixed strategies. Thus each individual can be represented by a vector $\mathbf{y} \in S_n$. The population is summarised by a distribution F on S_n . The mean strategy in the population is given by $\mathbf{x} = \int \mathbf{y} dF$ and the symmetric covariance matrix $Q_m = \int (\mathbf{x} - \mathbf{y})(\mathbf{x} - \mathbf{y}) dF$ (m is for mixed-strategy dynamic). Zeeman worked only with distributions that were *full*, that is, distributions for which Q_m has maximal rank amongst those populations having the same mean \mathbf{x} . As noted above, Zeeman justified this restriction by appealing to mutations. Summarising his results, we have

Lemma 5 *If \mathbf{x} is in the interior of S_n then $\mathbf{z} \cdot Q_m \mathbf{z} > 0$ for any \mathbf{z} which is not a multiple of \mathbf{u} . (Zeeman 1981, p265).*

Assuming as for the pure strategy replicator dynamic that the proportional growth rate of a strategy is equal to the difference between its and the average payoff gives

$$\dot{f}(\mathbf{y}) = f(\mathbf{y})[\mathbf{y} \cdot \mathbf{A}\mathbf{x} - \mathbf{x} \cdot \mathbf{A}\mathbf{x}]$$

and hence

Lemma 6 *The dynamic for the mean mixed strategy satisfies $\dot{\mathbf{x}} = Q_m \mathbf{A}\mathbf{x}$. (Zeeman 1981, p266).*

We can find similar results for the type of learning dynamics considered by Harley (1982), Börgers and Sarin (1993), Crawford (1989) and Roth and Erev (1995). This may seem strange in that, first, Börgers and Sarin rightly point out this learning process when aggregated across a population of players is not identical to the replicator dynamics for either pure or mixed strategies, and that, second, Crawford proves that in such a large population, under such dynamics the mixed-strategy equilibrium of a simple game like (2) is unstable. However, Crawford's definition of a mixed-strategy equilibrium is the state where every agent plays the equilibrium mixed-strategy, that is, in game (2), they all play their first strategy with probability a . However, I would argue that in a random-mixing population this definition is over-strict. It is possible to have a state where the average strategy in the population, and hence, the expected strategy of an opponent, is equal to the mixed strategy equilibrium, although no agent plays the exact mixed strategy equilibrium profile. For example, the i th member of the population could play her first strategy with probability $a + \epsilon_i$ with $\sum \epsilon_i = 0$.

We assume, as for fictitious play, that each player has a vector \mathbf{w} , each element representing the "confidence" placed on each strategy. However, rather than choosing the strategy with the highest weight, each player plays strategy i with probability

$$y_i = \frac{w_i}{\sum_{i=1}^n w_i} = \frac{w_i}{W}.$$

Thus, here, in a similar way to the model of Zeeman, we can represent each individual as a point $\mathbf{y} \in S_n$, distributed according to a function F . However, here we have to take account of the magnitude of W , the sum of an agents' weights. We assume that they are distributed on \mathbf{R} according to a function G , and let H be the joint distribution function (incorporating F and G) on $S_n \times \mathbf{R}$. And again, in a large random-mixing population, the probability of meeting an opponent playing strategy i will be x_i , where again we define the population mean as $\mathbf{x} = \int \mathbf{y} dF$. However, rather than strategy distributions being changed according to an evolutionary process, each individual learns by adjusting the probability

that she plays each strategy in relation to the payoff that the strategy earns. If a strategy is chosen, and playing that strategy yields a positive payoff, then the probability of playing that strategy is “reinforced” by the payoff earned. In particular, if an individual plays strategy i against an opponent playing strategy j , then the i th element of \mathbf{w} is increased by the resulting payoff

$$w_i(t + \Delta t) = w_i(t) + a_{ij}.$$

However, all other elements of \mathbf{w} remain unchanged. This is the “Basic Model” of Roth and Erev (1995), who give a number of reasons why this may be a reasonable approximation of human learning. Thus the expected change is given by,

$$E [w_i(t + \Delta t)] = y_i (w_i(t) + (A\mathbf{x})_i) + (1 - y_i)w_i(t). \quad (20)$$

There are three important differences between this learning rule and fictitious play. First, it is stochastic, not deterministic. Second, while under fictitious play, agents have a limited capacity for assessing what they might have received if they had used some other strategy, here agents only consider what actions they actually play and what payoffs they actually receive (this type of learning model was developed to analyse animal behaviour). Third, for the probabilities to remain well defined, we must require all payoffs to be non-negative¹⁰, and that all agents start with all elements of their vector \mathbf{w} strictly positive. From (20), we can obtain

$$E [y_i(t + \Delta t) - y_i(t)] = \frac{y_i ((A\mathbf{x})_i - \mathbf{y} \cdot A\mathbf{x})}{W + \mathbf{y} \cdot A\mathbf{x}}. \quad (21)$$

This is a special case¹¹ of the RPS rule of Harley(1982). Crawford (1989) characterises individual behaviour in a large population of players by the

¹⁰Either we consider only games with positive payoffs, or we add a positive constant to all payoffs sufficiently large to make them positive. Clearly such a transformation would make no difference to a game’s strategic properties, though, in a dynamic context it can change the rate of adjustment. See the discussion of discrete time processes in the next section.

¹¹The equation (21) can be obtained by setting what Harley calls the “memory factor” to 1, and taking the continuous time limit.

deterministic continuous time equation,

$$\dot{y}_i = y_i[(Ax)_i - \mathbf{y} \cdot Ax]. \quad (22)$$

Börgers and Sarin (1993) show that by using a slightly different specification of the updating rule one can obtain a continuous time limit similar to Crawford's equation (22)¹². The advantage of the approach of Börgers and Sarin and Crawford is that learning behaviour is easier to characterise, but only at the cost of additional assumptions.

In any case, the next step is to derive an expression for the evolution of the population mean. If we think of the change made by each agent as a draw from the distribution that describes the population, $x_i(t+\Delta t) - x_i(t)$ is then the sample mean. Hence, the variance of the change in x_i is decreasing in the number of agents. Thus, if the population is infinite, then the evolution of the population mean will be deterministic (the case of a finite population will be dealt with in the next section). We have

$$\begin{aligned} x_i(t + \Delta t) - x_i(t) &= \int E[y_i(t + \Delta t) - y_i(t)] dH \\ &= \int y_i[(Ax)_i - \mathbf{y} \cdot Ax]/(W + \mathbf{y} \cdot Ax) dH \\ &= \int y_i[\mathbf{e}_i - \mathbf{y}]/(W + \mathbf{y} \cdot Ax) dH \cdot Ax. \end{aligned}$$

where \mathbf{e}_i is a vector of zeros except for a 1 in the i th position and $W + \mathbf{y} \cdot Ax > 0$ (by the assumption of non-negative payoffs). We take the continuous time limit. This in turn gives us,

$$\dot{\mathbf{x}} = Q_g Ax \quad (23)$$

where the g -subscript is for gradient learning. The diagonal of Q_g has the form $\int y_i(1 - y_i)/(W + \mathbf{y} \cdot Ax) dH$, the off-diagonal $-\int y_i y_j/(W + \mathbf{y} \cdot Ax) dH$. Hence Q_g is symmetric and $Q_g \mathbf{u} = \mathbf{0}$. Clearly $\mathbf{z} \cdot Q_g \mathbf{z} = \sum_{i \neq j} \int y_i y_j/(W + \mathbf{y} \cdot Ax) dH (z_i - z_j)^2 \geq 0$. Consequently Q_g is positive semi-definite. To obtain the model of either Börgers and Sarin (1993) or Crawford(1989) it simply necessary to set $W + \mathbf{y} \cdot Ax = 1$ for all agents. Clearly this would not change the conclusion that although $Q_g \neq Q_m$,

¹²It would be the same if Börgers and Sarin considered as did Crawford a single random-mixing population.

Proposition 7 *The mean of the mixed strategy replicator dynamic and the mean of the gradient learning process are positive definite dynamics.*

This, together with Proposition 5, extends the existing results on gradient dynamics.

An Example. Take the game (2), assume $a = .5$, that $F(y_1) = y_1^2$, and hence $x_1 = 2/3$. Under the mixed strategy replicator dynamics, we have $\dot{f}(y_1) = 2y_1[1/9 - y_1/3]$. That is, those agents playing the first strategy with probability less than one third, and hence far from the equilibrium strategy, are increasing in number. For the gradient dynamics, we have $\dot{y}_1 = -y_1(1 - y_1)/(6W + 2 - y_1)$. In words, all agents are decreasing the weight they place on their first strategy. This also demonstrates the difference between the two dynamics. The evolutionary dynamic replaces badly-performing agents by better performers¹³, under the gradient dynamics, all agents respond to the situation by changing strategy. As Crawford (1989) discovered, the state where all agents have $y_1 = 0.5$ is not going to be stable. In this example, the agents who are currently playing the “equilibrium” mixed strategy ($y_1 = 0.5$) are respectively dying off and moving away from it. However, for both dynamics we have $\dot{x}_1 = Q_{11} [1/2 - x_1]$, and hence the mean strategy clearly approaches the equilibrium¹⁴.

7 A Stochastic Model

It is clear that the assumptions of continuous time and an infinite population have smoothed the path toward the results that we have obtained. However, how much can be maintained without them? This matters

¹³Though perhaps this type of dynamic could be reproduced in a population that learns by imitation.

¹⁴Harley (1982, p624) reproduces two graphs of the results he obtained from simulations of a similar game using his learning model. Two things are apparent: the population mean approaches the mixed strategy equilibrium, the strategy of individual players (typically) does not.

if we want, for example, to use this type of model, as do Gale et al. (1995) and Roth and Erev (1995), to predict the outcome of experimental games. It is possible to recreate our assumption of random-matching in an experimental environment. Perhaps it is possible to repeat the matching sufficiently rapidly to approximate continuous time. However, a typical research budget is not adequate to support an infinite number of experimental subjects.

First, let us consider learning dynamics in discrete time. Consider a positive definite dynamic such that

$$\mathbf{x}(t+1) = \mathbf{x}(t) + QA\mathbf{x}, \quad (24)$$

where Q again satisfies the five conditions outlined above. In this case, pure strategies which are regular ESSs will be asymptotically stable, the second part of the proof of Proposition 5 applying equally well in discrete time. The problem is, as always, with mixed strategies. From (24), the linearisation at a fully mixed fixed point \mathbf{q} will be

$$I + Q(\mathbf{q})A. \quad (25)$$

As we have shown, the eigenvalues of QA are negative. If however, they are too “large”, the absolute values of the eigenvalues of $I + QA$ will be greater than one. So it is possible for a discrete time positive definite process to diverge from a mixed ESS. This is going to depend on the magnitude of the change in strategy distribution made each period. In the case of a pure strategy equilibrium, it must be true that $\|\mathbf{x} - \mathbf{q}\| \ll \|QA\mathbf{x}\|$ otherwise the dynamic would jump over the fixed point and out of the simplex. In contrast, unless the rate of change is sufficiently slow, it is possible to shoot right past a mixed-strategy equilibrium. Note that, for example, for the discrete time replicator dynamics given in (3), the rate of adjustment is decreasing in the constant D . Hence, stability of ESSs can be assured if D is sufficiently large. In the case of gradient learning, the rate of change is decreasing over time as the size of individuals’ weights (W in the notation of the last section) increases.

We now examine the relationship between the results obtained in Section 5 in a deterministic setting and those obtainable when the evolution of play is stochastic. Learning and evolutionary systems with noise have been studied intensively in the recent literature, for example by Canning (1992), Kandori et al. (1993), Young (1993), Binmore et al. (1993). However, while these models are driven by mistakes or mutations, the noise in this model presented in this section is a consequence of the matching mechanism and the consequential heterogeneity of information. We assume a population of N players repeatedly (we continue to work in discrete time) randomly matched according to Story 3. Under the decentralised matching scheme, Story 3, when the population of players is finite, each player may have a different estimate of the frequencies of strategies in the population according to his random history of opponents. We show, using the PFP process, that this produces significantly different results than the noise produced by mistakes.

In this context, gradient learning and fictitious play part company. A state \mathbf{q} is absorbing if $\Pr(\mathbf{x}(t+1) \neq \mathbf{q} \mid \mathbf{x}(t) = \mathbf{q}) = 0$. That is, if the system reaches an absorbing state, it will never leave. For gradient learning the vertices of the simplex are absorbing: if an agent plays any strategy with probability one, she can learn no new strategies. This is not the case for fictitious play. In a deterministic setting (Section 4), there was a one-to-one relationship between fixed points and Nash equilibria, but now only a subset of Nash equilibria are absorbing.

Proposition 8 *A state \mathbf{q} is absorbing for the stochastic PFP if and only if for any strategy i in the support of \mathbf{q} ,*

1. $a_{ii} \geq a_{ji} \forall j$ and
2. $a_{ii} \leq a_{ki} \forall k$ in the support of \mathbf{q} .

That is, an absorbing state must be a Nash equilibrium in pure strategies or an element of a continuum of Nash equilibria between two or more pure Nash equilibria.

Proof: If there is only one strategy i in the support of \mathbf{q} , then clearly Condition 1, which is also the condition for i to be a Nash equilibrium strategy, is necessary and sufficient for a state to be absorbing. This follows as if all agents play i , they must prefer it. Further encounters with opponents playing i , since it is a best reply to itself, can only reinforce that preference. If Condition 1 fails, and hence there exists another strategy j which is a better reply to i , then for all players $w_j - w_i$ will increase monotonically and at a constant rate. Within finite time, at least one agent will change strategy. If there is more than one strategy in the support of \mathbf{q} , if Conditions 1 and 2 hold, no sequence of matches will change any player's relative preference over strategies either in or not in the support of \mathbf{q} . However, for any agent playing strategy k (in the support of \mathbf{q}) there is a non-zero probability of being matched m times in succession with opponents playing some other strategy i in the support of \mathbf{q} . Clearly, there is a finite m such that this player will switch to i , if Condition 2 fails, and to j if Condition 1 does not hold. \square

One implication of this result is that the stochastic PFP process selects against weakly-dominated strategies in a way that evolutionary or learning processes that depend on the aggregate state of the population, do not. In (26) there is a game taken from Samuelson (1994). The strategy T (weakly) dominates M . However, there is a continuum of Nash equilibria, where a weight up to one half is placed on M . In Samuelson (1994), agents respond to the expected return to the different strategies, that is, the current overall state of the population. The matching scheme is Story 2 in our terminology. Hence all the elements of the continuum are fixed points of the adjustment process. Samuelson shows that even the introduction of noise generated by mistakes will not eliminate the play of the weakly dominated strategy M . However, by the above proposition, the only absorbing state for the PFP process is the pure strategy equilibrium. This is because the noise produced by the matching process implied by Story 3 in a finite population is different than that generated by mistakes or mutations. In the case of (26), when there is a mixture of T and M in the population, there is always a possibility, by meeting sufficient times, an opponent playing strategy M , for

a player to switch to strategy B . This causes selection against strategy M .

| | | | | |
|-----|-----|-----|-----|------|
| | T | M | B | |
| T | 1 | 1 | 1 | (26) |
| M | 1 | 1 | 0 | |
| B | 0 | 2 | 0 | |

We turn now to the question of stability of equilibria. Obviously, it is not possible to obtain the same type of convergence result as we did in a deterministic setting. What we can achieve is a result on the time averages of the learning process. Without the assumption of an infinite population the mean of the gradient learning process in a finite population satisfies¹⁵,

$$E[\mathbf{x}(t+1) - \mathbf{x}(t)] = Q_g A \mathbf{x}. \quad (27)$$

We can expect the evolution of \mathbf{x} to be similar to the deterministic case, particularly as it is already an average, whose variance is decreasing in the size of the population.

For fictitious play, the problems are worse. For example, one implication of Proposition 8 is that mixed ESSs are not absorbing¹⁶. Worse, they may not even exist. For example, it is not easy to support a mixed strategy equilibrium of $(1/3, 1/3, 1/3)$ when N is not divisible by 3. In the case of the PFP for each agent the expected change in beliefs will be the same as the actual change in beliefs in the deterministic model. That is, from (7),

$$E[\mathbf{p}_i(t+1)] = \mathbf{p}_i(t) + [(\mathbf{A}\mathbf{x})_{j \neq i} - (\mathbf{A}\mathbf{x})_i].$$

If we maintain the Continuity Assumption, from (14) we have

$$E[x_i(t+1) - x_i(t)] \approx \sum_{j \neq i} [(\mathbf{A}\mathbf{x})_i - (\mathbf{A}\mathbf{x})_j] f_{ij}(\mathbf{0}). \quad (28)$$

¹⁵This is obtained simply by integrating (21) across the population.

¹⁶Mixed ESSs are always isolated and are never linear combinations of two pure strategy equilibria.

This approximation may be less than perfect. Hence, we assume

$$\mathbf{x}(t+1) - \mathbf{x}(t) = Q\mathbf{A}\mathbf{x} + o(\mathbf{x}). \quad (29)$$

This last object is a vector of random variables of length n , in which is wrapped up the errors generated by the approximation (28), the random element introduced by the matching scheme, and the inaccuracies introduced by representing a discrete state space (N players) by a continuous variable \mathbf{x} . We assume $o(\mathbf{x}) \cdot \mathbf{u} = 0$ (this is necessary simply to insure that the dynamic is well defined and remains within the simplex), and $E[o(\mathbf{x}) \cdot o(\mathbf{x})]$ is finite. That is, all the o_i have finite variance. Finally, we need $\lim_{x_i \rightarrow 0} o_i \geq 0$, again to ensure that the dynamic does not leave the simplex. However, it also reflects the fact that in a state where only one strategy is played, the noise caused by a random matching process is non-existent. We do not assume that $E[o(\mathbf{x})] = 0$, but the somewhat weaker condition that $\lim_{T \rightarrow \infty} \frac{1}{T} \sum o(\mathbf{x}) = 0$. That is, all the errors discussed are in the long-run unbiased. Note that (29) includes (27) as a special case.

Let

$$\mathbf{m}_{1T} = \frac{1}{T} \sum_{t=1}^T \mathbf{x}(t), \quad P_{1T} = \frac{1}{T} \sum_{t=1}^T Q(t).$$

That is, \mathbf{m}_{1T} is the time average of \mathbf{x} over the periods 1 to T . Then,

$$\mathbf{m}_{1T+1} = \mathbf{m}_{0T} + P\mathbf{A}\mathbf{m}_{0T} + \frac{1}{T+1} \sum_0^T o(\mathbf{x}) \quad (30)$$

If T is “large”, the last term in the above equation should tend to zero. In other words, as time goes by the behaviour of the time average \mathbf{m} of the discrete time stochastic process should approximate the evolution of the strategy frequencies \mathbf{x} under the deterministic discrete time process (24) ever more closely. As for the deterministic process, stability will depend on the rate of adjustment. Thus, we state the next result as a conjecture as we cannot prove that, even for an ESS, that given the linearisation (25), that all eigenvalues have absolute value less than one.

Conjecture 1 *If a game possesses an unique mixed ESS, \mathbf{q} , then under fictitious play or gradient learning, from any fully-mixed initial conditions, $\lim_{t \rightarrow \infty} \mathbf{m}_{1t} = \mathbf{q}$.*

This is on one level a very weak claim. With gradient learning dynamics, for example, even though a game possesses an unique mixed ESS, all we ask is that the time average of the average strategy should converge to the equilibrium. Thus we would not (necessarily) have convergence for either individual players or individual time periods. However, it is a result that can be tested by experiment, particularly in the sense that, as we will now see, there are games which possess an unique Nash equilibrium yet for which even this weak form of convergence will not hold.

Figure 2 graphs the time averages of play produced by simulations of the PFP process in a population size of 12. The game played was

$$A = \begin{array}{|c|c|c|} \hline 0 & a_1 & -b_1 \\ \hline -b_2 & 0 & a_2 \\ \hline a_3 & -b_3 & 0 \\ \hline \end{array} \quad a_i, b_i > 0, \quad i = 1, 2, 3 \quad (31)$$

which possesses an unique mixed equilibrium at $(1/3, 1/3, 1/3)$. Simulation (a) was produced with $a_i = 3, b_i = 1 \quad i = 1, 2, 3$, and (b) with $a_i = 1, b_i = 3$. In case (a), the equilibrium is negative definite and there is convergence, but in case (b), even the time averages diverge. We explain this result in the next section.

8 Games without ESSs

Since the concept of an ESS is a strong refinement on Nash equilibrium and consequently there are many games which do not possess any equilibrium which satisfies its conditions, one might wonder how positive definite dynamics perform in these cases. For any constant-sum game for any \mathbf{x} , $\mathbf{x} \cdot A\mathbf{x} = v$, where v is the value of the game. It follows, if the game has a fully mixed equilibrium \mathbf{q} , that $(\mathbf{x} - \mathbf{q}) \cdot A(\mathbf{x} - \mathbf{q}) = 0$. From Proposition 5 and in particular (18) we have that,





Figure 3: Time averages (\mathbf{m}_{1T}) of play from a simulation of fictitious play in a random mixing population.

Corollary 1 *The eigenvalues of any differentiable positive definite system at a fully mixed Nash equilibrium of a zero-sum game have zero real part.*

This result unfortunately is of the “anything can happen” type. For the linear dynamics (17), because they are linear, the Corollary implies that such an equilibrium must be a neutrally stable centre (it is easy to check that $V = \frac{1}{2}(\mathbf{x} - \mathbf{q}) \cdot (\mathbf{x} - \mathbf{q})$ is a constant of motion in this case). For non-linear dynamics the fact that their linearisations have zero eigenvalues may hide asymptotic stability or instability.

Secondly, there are games which possess equilibria which are positive definite. It is an obvious corollary of Proposition 5 that positive definite dynamics diverge from such equilibria. This can prove useful in terms of equilibrium selection. Unstable positive definite equilibria can be rejected in favour of stable ESSs. This works well in games with both ESSs and positive definite equilibria. But the game (31) has an unique equilibrium which, for example, for $a_i = 1, b_i = 3, i = 1, 2, 3$ is positive definite. Hence, no positive definite dynamic can converge. This might seem problematic, but in fact it offers a strong empirical prediction. For rational players under the full-information assumptions of conventional game theory, for a game with an unique Nash equilibrium it should not matter whether it is positive or negative definite. However, we can conjecture that in a random-matching environment under experimental conditions, the strategy frequencies of human subjects would converge if, for example, $a_i = 3$ and $b_i = 1$ but not if $a_i = 1$ and $b_i = 3$ (Figure 2 shows the results of simulations of fictitious play for these parameter values). This conjecture we can make with a degree of confidence because so many different specifications of adaptive learning are consistent with positive definite dynamics. Furthermore, in discrete time, if \mathbf{q} is a positive definite equilibrium, and hence QA has positive eigenvalues, then all the eigenvalues of the linearisation (25) are clearly greater than one and the equilibrium will most certainly be unstable. Such divergence is not necessarily “irrational” or “myopic”. Indeed, if $a_i = 1, b_i = 3, i = 1, 2, 3$

average payoffs are at a minimum at the mixed equilibrium. Divergence increases average payoffs.

The robustness of these results, however, does depend on the property of positive or negative definiteness. For equilibria which are neither positive neither negative definite, it is possible for stability properties to vary according to the exact specification of the dynamics. Such equilibria can be attractors or repellers. Using (31) again as an example, the pure strategy replicator dynamics converge iff $a_1 a_2 a_3 > b_1 b_2 b_3$, the linear dynamics iff $a_1 + a_2 + a_3 > b_1 + b_2 + b_3$, while simulation suggests that the PFP dynamics will converge to any equilibrium of the game which is not positive definite.

9 Conclusion

There has been some debate as to whether the replicator dynamics, in spite of their biological origins, can serve as a learning dynamic for human populations. The results obtained here on one level give some support to the skeptics. The aggregation of learning behaviour across a large population is not in general identical to the replicator dynamics, in either their pure or mixed strategy formulation. However, it is clear that all these dynamics, whether of learning or evolution, share many of the same properties.

This is valuable in that, as the literature on learning and evolution has been growing at a significant rate over the past few years, there has been a proliferation of different models and consequently different results. The hope here is that we have obtained a result that is reasonably robust: ESSs are asymptotically stable for many apparently different adaptive processes when these processes are aggregated across a large random-mixing population. An ESS is quite a strong refinement on Nash equilibrium. Furthermore, it has been discredited in the eyes of some because it does not correspond exactly to asymptotic stability under the pure strategy replicator dynamics (Proposition 1). However,

these are not the only dynamics of interest, and for results on stability that are robust to different specifications, the concept of ESS is the one that is relevant. In extending existing results on fictitious play, gradient learning and mixed-strategy replicator dynamics, it has been the negative definiteness of ESSs which has been essential.

Researchers have begun to test the predictions of models of learning and evolution by carrying out experiments. The results presented in this paper may be relevant in several ways. First, Mookherjee and Sopher (1994), for example, attempt to determine whether fictitious play or gradient type rules best describe the learning behaviour of their subjects. As we have shown, the differences between these two types of model, in a random-matching environment at least, are smaller than previously thought. Our results would also point to a reason why Gale et al. (1995), using the replicator dynamics, and Roth and Erev (1995), using a gradient type learning process obtain similar results in trying to simulate the behaviour of experimental subjects playing the ultimatum bargaining game. Second, there has been some debate (Brown and Rosenthal, 1990; Binmore, Swierzbinski, and Proulx, 1994) about what constitutes convergence to equilibrium in experimental games. What we show here is that it may be foolish to expect more than convergence in the average strategy in a population of players. Thirdly, we offer further predictions to be tested. Games which possess ESSs should converge. For games which possess positive definite equilibria, our predictions are equally clear. Learning processes should not converge to such equilibria.

Finally, as we noted in Section 1, under fictitious play for some mixed strategy equilibria there is convergence in beliefs without convergence in play. In the random-mixing models considered here, the opposite is possible. The distribution of strategies in the population matches exactly the equilibrium strategy profile. However, individual agents play any mix over the strategies in its support, including a single pure strategy. One might say that none has "learnt" the mixed strategy equilibrium, but equally, given the assumption of random matching none has an incentive to change strategy.

References

- Binmore, K., Samuelson L.** (1994) "Muddling through: noisy equilibrium selection," Working Paper, University College, London; University of Wisconsin.
- Binmore, K., Samuelson L., Vaughan R.** (1993) "Musical chairs: modelling noisy evolution," Working Paper, University College, London; University of Wisconsin.
- Binmore, K., Swierzbinski, J., Proulx, C.** (1994) "Does minimax work? An experimental study," Working Paper. University College, London.
- Börgers, T., Sarin R.** (1993) "Learning through reinforcement and replicator dynamics," Working Paper, University College, London.
- Boylan, R.T.** (1992) "Laws of large numbers for dynamical systems with randomly matched individuals," *J. Econ. Theory*, **57**, 473-506.
- Brown, G.W.** (1951). "Iterative solutions of fictitious play," in *Activity Analysis of Production and Allocation* (T.C. Koopmans, Ed.). New York: Wiley.
- Brown, J., Rosenthal, R.** (1990) "Testing the minimax hypothesis: a re-examination of O'Neill's game experiment," *Econometrica*, **58**, 1065-1081.
- Canning, D.** (1992). "Average behaviour in learning models," *J. Econ. Theory*, **57**, 442-472.
- Crawford, V.** (1985). "Learning behaviour and mixed-strategy Nash equilibria," *J. Econ. Behav. Organ.*, **6**, 69-78.
- Crawford, V.** (1989) "Learning and mixed strategy equilibria in evolutionary games," *J. Theor. Biol.*, **140**, 537-550.
- Friedman, D.** (1991) "Evolutionary games in economics," *Econometrica*, **59**, 637-666.

- Fudenberg, D., Kreps D.** (1993). "Learning mixed equilibria," *Games Econ. Behav.*, **5**, 320-367.
- Fudenberg, D., Levine D.** (1993). "Steady state learning and Nash equilibrium," *Econometrica*, **61**, 547-573.
- Gale, J., Binmore, K., Samuleson, L.** (1995) "Learning to be imperfect: the ultimatum game," *Games Econ. Behav.*, **8**, 56-90.
- Harley, C.** (1981). "Learning the evolutionary stable strategy," *J. Theor. Biol.*, **89**, 611-633.
- Hines, W.G.S.** (1980). "Three characterisations of population strategy stability," *J. Appl. Prob.*, **17**, 333-340.
- Hofbauer, J., Sigmund K.** (1988). *The Theory of Evolution and Dynamical Systems*. Cambridge: Cambridge University Press.
- Jordan, J.** (1993). "Three problems in learning mixed strategy equilibria," *Games Econ. Behav.*, **5**, 368-386.
- Kalai, E., Lehrer E.** (1993). "Rational learning leads to Nash equilibrium," *Econometrica*, **61**, 1019-1045.
- Kandori, M., Mailath G. and Rob R.** (1993). "Learning, mutation and long run equilibrium in games," *Econometrica*, **61**, 29-56.
- Maynard Smith, J.** (1982). *Evolution and the Theory of Games*. Cambridge: Cambridge University Press.
- Milgrom, P., Roberts J.** (1991). "Adaptive and sophisticated learning in normal form games," *Games Econ. Behav.*, **3**, 82-100.
- Miyasawa, K.** (1961) "On the convergence of the learning process in a 2×2 non-zero sum game," Mimeo, Princeton University.
- Monderer, D., Shapley L.** (1993). "Fictitious play property for games with identical interests," forthcoming *J. Econ. Theory*.
- Mookherjee, D., Sopher, B.** (1994). "An experimental study of learning and decision costs in constant-sum games," working paper.

- Nachbar, J.H.** (1990). "Evolutionary selection dynamics in games: convergence and limit properties," *Int. J. Game Theory*, **19**, 59-89.
- Robinson, J.** (1951). "An Iterative Method of Solving a Game," *Ann. Math.*, **54**, 296-301.
- Roth, A.E., Erev, I.** (1995). "Learning in extensive-form games: experimental data and simple dynamic models in the intermediate term," *Games Econ. Behav.*, **8**, 164-212.
- Samuelson, L.** (1994). "Stochastic stability in games with alternative best replies," *J. Econ. Theory*, **64**, 35-65.
- Samuelson, L., Zhang J.** (1992). "Evolutionary stability in asymmetric games," *J. Econ. Theory*, **57**, 363-392.
- Shapley, L.** (1964) "Some topics in two person games," in eds. M. Dresher et al., *Advances in Game Theory*. Princeton: Princeton University Press.
- Sugden, R.** (1989) "Spontaneous order," *J. Econ. Perspectives*, **3.4**, 85-97.
- van Damme, E.** (1991) *The Perfection and Stability of Nash Equilibrium*. Second Edition. Berlin: Springer Verlag.
- Young, H.P.** (1993) "Conventional equilibria," *Econometrica*, **61**, 57-84.
- Zeeman, E.C.** (1981) "Dynamics of the evolution of animal conflicts," *J. Theor. Biol.*, **89**, 249-270.



EUI WORKING PAPERS

EUI Working Papers are published and distributed by the
European University Institute, Florence

Copies can be obtained free of charge
– depending on the availability of stocks – from:

The Publications Officer
European University Institute
Badia Fiesolana
I-50016 San Domenico di Fiesole (FI)
Italy

Please use order form overleaf



Publications of the European University Institute

Department of Economics Working Paper Series

To Department of Economics WP
 European University Institute
 Badia Fiesolana
 I-50016 San Domenico di Fiesole (FI)
 E-mail: publish@datacomm.iue.it
 Italy

From Name

Address

.....

.....

.....

(Please print)

- Please enter/confirm my name on EUI Economics Dept. Mailing List
- Please send me a complete list of EUI Working Papers
- Please send me a complete list of EUI book publications
- Please send me the EUI brochure Academic Year 1995/96

Please send me the following EUI ECO Working Paper(s):

No, Author

Title:

No, Author

Title:

No, Author

Title:

No, Author

Title:

Date Signature

**Working Papers of the Department of Economics
Published since 1993**

- ECO No. 93/1**
Carlo GRILLENZONI
Forecasting Unstable and Non-Stationary
Time Series
- ECO No. 93/2**
Carlo GRILLENZONI
Multilinear Models for Nonlinear Time
Series
- ECO No. 93/3**
Ronald M. HARSTAD/Louis PHILIPS
Futures Market Contracting When You
Don't Know Who the Optimists Are
- ECO No. 93/4**
Alan KIRMAN/Louis PHILIPS
Empirical Studies of Product Markets
- ECO No. 93/5**
Grayham E. MIZON
Empirical Analysis of Time Series:
Illustrations with Simulated Data
- ECO No. 93/6**
Tilman EHRBECK
Optimally Combining Individual
Forecasts From Panel Data
- ECO NO. 93/7**
V́ctor GÓMEZ/Agustín MARAVALL
Initializing the Kalman Filter with
Incompletely Specified Initial Conditions
- ECO No. 93/8**
Frederic PALOMINO
Informed Speculation: Small Markets
Against Large Markets
- ECO NO. 93/9**
Stephen MARTIN
Beyond Prices Versus Quantities
- ECO No. 93/10**
José María LABEAGA/Angel LÓPEZ
A Flexible Demand System and VAT
Simulations from Spanish Microdata
- ECO No. 93/11**
Maozu LU/Grayham E. MIZON
The Encompassing Principle and
Specification Tests
- ECO No. 93/12**
Louis PHILIPS/Peter MØLLGAARD
Oil Stocks as a Squeeze Preventing
Mechanism: Is Self-Regulation Possible?
- ECO No. 93/13**
Pieter HASEKAMP
Disinflation Policy and Credibility: The
Role of Conventions
- ECO No. 93/14**
Louis PHILIPS
Price Leadership and Conscious
Parallelism: A Survey
- ECO No. 93/15**
Agustín MARAVALL
Short-Term Analysis of Macroeconomic
Time Series *
- ECO No. 93/16**
Philip Hans FRANSES/Niels
HALDRUP
The Effects of Additive Outliers on Tests
for Unit Roots and Cointegration
- ECO No. 93/17**
Fabio CANOVA/Jane MARRINAN
Predicting Excess Returns in Financial
Markets
- ECO No. 93/18**
Iñigo HERGUERA
Exchange Rate Fluctuations, Market
Structure and the Pass-through
Relationship
- ECO No. 93/19**
Agustín MARAVALL
Use and Misuse of Unobserved
Components in Economic Forecasting
- ECO No. 93/20**
Torben HOLVAD/Jens Leth
HOUGAARD
Measuring Technical Input Efficiency for
Similar Production Units:
A Survey of the Non-Parametric
Approach

*out of print

ECO No. 93/21

Stephen MARTIN/Louis PHILIPS
Product Differentiation, Market Structure
and Exchange Rate Passthrough

ECO No 93/22

F. CANOVA/M. FINN/A. R. PAGAN
Evaluating a Real Business Cycle Model

ECO No 93/23

Fabio CANOVA
Statistical Inference in Calibrated Models

ECO No 93/24

Gilles TEYSSIÈRE
Matching Processes in the Labour Market
in Marseilles. An Econometric Study

ECO No 93/25

Fabio CANOVA
Sources and Propagation of International
Business Cycles: Common Shocks or
Transmission?

ECO No. 93/26

Marco BECHT/Carlos RAMÍREZ
Financial Capitalism in Pre-World War I
Germany: The Role of the Universal
Banks in the Financing of German
Mining Companies 1906-1912

ECO No. 93/27

Isabelle MARET
Two Parametric Models of Demand,
Structure of Market Demand from
Heterogeneity

ECO No. 93/28

Stephen MARTIN
Vertical Product Differentiation, Intra-
industry Trade, and Infant Industry
Protection

ECO No. 93/29

J. Humberto LOPEZ
Testing for Unit Roots with the k-th
Autocorrelation Coefficient

ECO No. 93/30

Paola VALBONESI
Modelling Interactions Between State and
Private Sector in a "Previously" Centrally
Planned Economy

ECO No. 93/31

Enrique ALBEROLA ILA/J. Humberto
LOPEZ/Vicente ORTOS RIOS
An Application of the Kalman Filter to
the Spanish Experience in a Target Zone
(1989-92)

ECO No. 93/32

Fabio CANOVA/Morten O. RAVN
International Consumption Risk Sharing

ECO No. 93/33

Morten Overgaard RAVN
International Business Cycles: How
much can Standard Theory Account for?

ECO No. 93/34

Agustín MARAVALL
Unobserved Components in Economic
Time Series *

ECO No. 93/35

Sheila MARNIE/John
MICKLEWRIGHT
Poverty in Pre-Reform Uzbekistan:
What do Official Data Really Reveal? *

ECO No. 93/36

Torben HOLVAD/Jens Leth
HOUGAARD
Measuring Technical Input Efficiency for
Similar Production Units:
80 Danish Hospitals

ECO No. 93/37

Grayham E. MIZON
A Simple Message for Autocorrelation
Correctors: DON'T

ECO No. 93/38

Barbara BOEHNLEIN
The Impact of Product Differentiation on
Collusive Equilibria and Multimarket
Contact

ECO No. 93/39

H. Peter MØLLGAARD
Bargaining and Efficiency in a
Speculative Forward Market

- ECO No. 94/1**
Robert WALDMANN
Cooperatives With Privately Optimal
Price Indexed Debt Increase Membership
When Demand Increases
- ECO No. 94/2**
Tilman EHRBECK/Robert
WALDMANN
Can Forecasters' Motives Explain
Rejection of the Rational Expectations
Hypothesis?
- ECO No. 94/3**
Alessandra PELLONI
Public Policy in a Two Sector Model of
Endogenous Growth *
- ECO No. 94/4**
David F. HENDRY
On the Interactions of Unit Roots and
Exogeneity
- ECO No. 94/5**
Bernadette GOVAERTS/David F.
HENDRY/Jean-François RICHARD
Encompassing in Stationary Linear
Dynamic Models
- ECO No. 94/6**
Luigi ERMINI/Dongkoo CHANG
Testing the Joint Hypothesis of Rational-
ity and Neutrality under Seasonal Coin-
tegration: The Case of Korea
- ECO No. 94/7**
Gabriele FIORENTINI/Agustín
MARAVALL
Unobserved Components in ARCH
Models: An Application to Seasonal
Adjustment *
- ECO No. 94/8**
Niels HALDRUP/Mark SALMON
Polynomially Cointegrated Systems and
their Representations: A Synthesis
- ECO No. 94/9**
Mariusz TAMBORSKI
Currency Option Pricing with Stochastic
Interest Rates and Transaction Costs:
A Theoretical Model
- ECO No. 94/10**
Mariusz TAMBORSKI
Are Standard Deviations Implied in
Currency Option Prices Good Predictors
of Future Exchange Rate Volatility?
- ECO No. 94/11**
John MICKLEWRIGHT/Gyula NAGY
How Does the Hungarian Unemploy-
ment Insurance System Really Work? *
- ECO No. 94/12**
Frank CRITCHLEY/Paul
MARRIOTT/Mark SALMON
An Elementary Account of Amari's
Expected Geometry
- ECO No. 94/13**
Domenico Junior MARCHETTI
Procyclical Productivity, Externalities
and Labor Hoarding: A Reexamination of
Evidence from U.S. Manufacturing
- ECO No. 94/14**
Giovanni NERO
A Structural Model of Intra-European
Airline Competition
- ECO No. 94/15**
Stephen MARTIN
Oligopoly Limit Pricing: Strategic
Substitutes, Strategic Complements
- ECO No. 94/16**
Ed HOPKINS
Learning and Evolution in a
Heterogeneous Population
- ECO No. 94/17**
Berthold HERRENDORF
Seigniorage, Optimal Taxation, and Time
Consistency: A Review
- ECO No. 94/18**
Frederic PALOMINO
Noise Trading in Small Markets *
- ECO No. 94/19**
Alexander SCHRADER
Vertical Foreclosure, Tax Spinning and
Oil Taxation in Oligopoly
- ECO No. 94/20**
Andrzej BANIAK/Louis PHLIPS
La Pléiade and Exchange Rate Pass-
Through
- ECO No. 94/21**
Mark SALMON
Bounded Rationality and Learning;
Procedural Learning

*out of print

ECO No. 94/22

Isabelle MARET
Heterogeneity and Dynamics of
Temporary Equilibria: Short-Run Versus
Long-Run Stability

ECO No. 94/23

Nikolaos GEORGANTZIS
Short-Run and Long-Run Cournot
Equilibria in Multiproduct Industries

ECO No. 94/24

Alexander SCHRADER
Vertical Mergers and Market Foreclosure:
Comment

ECO No. 94/25

Jeroen HINLOOPEN
Subsiding Cooperative and Non-
Cooperative R&D in Duopoly with
Spillovers

ECO No. 94/26

Debora DI GIOACCHINO
The Evolution of Cooperation:
Robustness to Mistakes and Mutation

ECO No. 94/27

Kristina KOSTIAL
The Role of the Signal-Noise Ratio in
Cointegrated Systems

ECO No. 94/28

Agustín MARAVALL/Víctor GÓMEZ
Program SEATS "Signal Extraction in
ARIMA Time Series" - Instructions for
the User

ECO No. 94/29

Luigi ERMINI
A Discrete-Time Consumption-CAP
Model under Durability of Goods, Habit
Formation and Temporal Aggregation

ECO No. 94/30

Debora DI GIOACCHINO
Learning to Drink Beer by Mistake

ECO No. 94/31

Víctor GÓMEZ/Agustín MARAVALL
Program TRAMO "Time Series
Regression with ARIMA Noise, Missing
Observations, and Outliers" -
Instructions for the User

ECO No. 94/32

Ákos VALENTINYI
How Financial Development and
Inflation may Affect Growth

ECO No. 94/33

Stephen MARTIN
European Community Food Processing
Industries

ECO No. 94/34

Agustín MARAVALL/Christophe
PLANAS
Estimation Error and the Specification of
Unobserved Component Models

ECO No. 94/35

Robbin HERRING
The "Divergent Beliefs" Hypothesis and
the "Contract Zone" in Final Offer
Arbitration

ECO No. 94/36

Robbin HERRING
Hiring Quality Labour

ECO No. 94/37

Angel J. UBIDE
Is there Consumption Risk Sharing in the
EEC?

ECO No. 94/38

Berthold HERRENDORF
Credible Purchases of Credibility
Through Exchange Rate Pegging:
An Optimal Taxation Framework

ECO No. 94/39

Enrique ALBEROLA ILA
How Long Can a Honeymoon Last?
Institutional and Fundamental Beliefs in
the Collapse of a Target Zone

ECO No. 94/40

Robert WALDMANN
Inequality, Economic Growth and the
Debt Crisis

ECO No. 94/41

John MICKLEWRIGHT/
Gyula NAGY
Flows to and from Insured
Unemployment in Hungary

ECO No. 94/42

Barbara BOEHNLEIN
The Soda-ash Market in Europe:
Collusive and Competitive Equilibria
With and Without Foreign Entry

ECO No. 94/43

Hans-Theo NORMANN
Stackelberg Warfare as an Equilibrium
Choice in a Game with Reputation Effects

ECO No. 94/44

Giorgio CALZOLARI/Gabriele
FIORENTINI
Conditional Heteroskedasticity in
Nonlinear Simultaneous Equations

ECO No. 94/45

Frank CRITCHLEY/Paul MARRIOTT/
Mark SALMON
On the Differential Geometry of the Wald
Test with Nonlinear Restrictions

ECO No. 94/46

Renzo G. AVESANI/Giampiero M.
GALLO/Mark SALMON
On the Evolution of Credibility and
Flexible Exchange Rate Target Zones

ECO No. 95/1

Paul PEZANIS-CHRISTOU
Experimental Results in Asymmetric
Auctions - The 'Low-Ball' Effect

ECO No. 95/2

Jeroen HINLOOPEN/Rien
WAGENVOORT
Robust Estimation: An Example

ECO No. 95/3

Giampiero M. GALLO/Barbara PACINI
Risk-related Asymmetries in Foreign
Exchange Markets

ECO No. 95/4

Santanu ROY/Rien WAGENVOORT
Risk Preference and Indirect Utility in
Portfolio Choice Problems

ECO No. 95/5

Giovanni NERO
Third Package and Noncooperative
Collusion in the European Airline
Industry

ECO No. 95/6

Renzo G. AVESANI/Giampiero M.
GALLO/Mark SALMON
On the Nature of Commitment in Flexible
Target Zones and the Measurement of
Credibility: The 1993 ERM Crisis

ECO No. 95/7

John MICKLEWRIGHT/Gyula NAGY
Unemployment Insurance and Incentives
in Hungary

ECO No. 95/8

Kristina KOSTIAL
The Fully Modified OLS Estimator as a
System Estimator: A Monte-Carlo
Analysis

ECO No. 95/9

Günther REHME
Redistribution, Wealth Tax Competition
and Capital Flight in Growing
Economies

ECO No. 95/10

Grayham E. MIZON
Progressive Modelling of
Macroeconomic Time Series: The LSE
Methodology

ECO No. 95/11

Pierre CAHUC/Hubert KEMPF
Alternative Time Patterns of Decisions
and Dynamic Strategic Interactions

ECO No. 95/12

Tito BOERI
Is Job Turnover Countercyclical?

ECO No. 95/13

Luisa ZANFORLIN
Growth Effects from Trade and
Technology

ECO No. 95/14

Miguel JIMÉNEZ/Domenico
MARCHETTI, jr.
Thick-Market Externalities in U.S.
Manufacturing: A Dynamic Study with
Panel Data

ECO No. 95/15

Berthold HERRENDORF
Exchange Rate Pegging, Transparency,
and Imports of Credibility

*out of print

ECO No. 95/16

Günther REHME

**Redistribution, Income cum Investment
Subsidy Tax Competition and Capital
Flight in Growing Economies**

ECO No. 95/17

Tito BOERI/Stefano SCARPETTA

**Regional Dimensions of Unemployment
in Central and Eastern Europe and Social
Barriers to Restructuring**

ECO No. 95/18

Bernhard WINKLER

**Reputation for EMU - An Economic
Defence of the Maastricht Criteria**

ECO No. 95/19

Ed HOPKINS

Learning, Matching and Aggregation

